



HAL
open science

o-minimal expansions of real closed fields and completeness in the sense of Scott

Olivier Frécon

► **To cite this version:**

Olivier Frécon. o-minimal expansions of real closed fields and completeness in the sense of Scott. 2015. hal-03704341

HAL Id: hal-03704341

<https://univ-poitiers.hal.science/hal-03704341v1>

Preprint submitted on 24 Jun 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

O-MINIMAL EXPANSIONS OF REAL CLOSED FIELDS AND COMPLETENESS IN THE SENSE OF SCOTT

OLIVIER FRÉCON

ABSTRACT. We consider an o -minimal expansion $\mathcal{M}_0 = (R_0, <, +, \dots)$ of a real closed field, and a real closed field R , complete in the sense of D. Scott, containing R_0 as a dense subfield. We show that \mathcal{M}_0 has an elementary extension $\mathcal{M} = (R, <, +, \dots)$ with domain R . Moreover, such a structure \mathcal{M} with domain R is unique.

NOTE

In an unpublished article, Antonigliulo Fornasiero proved a more general result than the main theorem of this paper. Indeed, he showed a similar result for d -minimal expansions of a real closed field [1, Proposition 11.6]. However our proofs are very different.

1. INTRODUCTION

By the Compactness Theorem, it is easy to elementarily embed any expansion $\mathcal{M}_0 = (R_0, <, +, \dots)$ of a real closed field in an expansion $\mathcal{M} = (R, <, +, \dots)$ of a *Scott-complete* real closed field, that is complete in the sense of Dana Scott (Definition 1.1). However, we have little control on the size of the elementary extension obtained. For instance, if R_0 is countable, it is possible that the field R has no countable dense subfield. The main result of this paper shows that any o -minimal expansion of a real closed field R_0 is elementarily embedded in an expansion of a Scott-complete real closed field, in which R_0 is dense (Theorem 1.2). We note that, since we consider an expansion of a real closed field, not just the field structure, the model completeness of the theory of real closed fields will not help us.

We recall the definition of a *complete* real closed field in the sense of [2].

Definition 1.1. – *If K and L are ordered fields and $K \subseteq L$, then K is dense in L if between any two distinct elements of L there lies an element of K .*

A given ordered field is called Scott-complete if it has no proper extension to an ordered field in which the given field is dense.

The main result of this paper is the following.

Theorem 1.2. – *Any o -minimal expansion of a real closed field R_0 has an elementary extension of domain a Scott-complete field R in which R_0 is dense. Moreover, for a fixed field R , this elementary extension is unique.*

Date: May 4, 2015.

2010 Mathematics Subject Classification. 03C64.

Key words and phrases. Real closed field, o -minimal structure.

2. NOTATIONS AND PRELIMINARY RESULTS

For the rest of this paper, we fix an \mathcal{o} -minimal expansion $\mathcal{M}_1 = (R_0, <, \dots)$ of a real closed field R_0 , and we denote by R a Scott-complete field in which R_0 is dense: its existence is ensured by Fact 2.1 below.

Fact 2.1. – [2, Theorem 1] *Given any ordered field K , there is a Scott-complete ordered field \hat{K} in which K is dense. Any other Scott-complete ordered field in which K is dense is isomorphic to \hat{K} by a unique isomorphism that is the identity on K .*

In the following, we say that a set X is *definable* if it is definable in the structure \mathcal{M}_1 .

For each integer k and each subset A of R^k , we denote by \bar{A} its (topological) closure in R^k . For each subset X of R_0^k , we denote by \check{X} the union of subsets F for F a R_0 -closed definable subset contained in X .

For clarity, we use a very different notation for the closure in R_0^k of a subset X of R_0^k : we denote it by $\text{cl } X$. Moreover, we denote by ∂X its frontier: $\partial X = \text{cl } X \setminus X$.

For any element $x = (x_1, \dots, x_k)$ of R^k , we consider

$$|x| = \max\{|x_i| \mid i \in \{1, \dots, k\}\}$$

Remark 2.2. –

- For any subset X of R_0^k , it follows from the definition of \check{X} that $X = \check{X} \cap R_0^k$.
- The proof of Theorem 1.2 will show that the elementary extension of \mathcal{M}_1 of domain R has the following property:
for any two integers k and m , if A is any subset of R_0^k defined by a formula $\varphi(\bar{x}, \bar{a})$ with free variables $\bar{x} = (x_1, \dots, x_k)$ and parameters $\bar{a} \in R_0^m$, then the subset of R^k defined by $\varphi(\bar{x}, \bar{a})$ is \bar{A} .

The first lemma is certainly well-known, but we could not find a reference for it.

Lemma 2.3. – *Let k be an integer, X a subset of R_0^k , and $f : X \rightarrow R_0$ be a uniformly continuous map. Then for each $x \in \bar{X}$, the following limit exists:*

$$\lim_{\substack{y \in X \\ y \rightarrow x}} f(y)$$

PROOF – Let $\varepsilon \in R_0^{>0}$. Since f is uniformly continuous on X , then we may associate with ε some $\delta(\varepsilon) \in R_0^{>0}$ such that whenever $|v - u| < \delta(\varepsilon)$ for u and v in X , we have $|f(v) - f(u)| < \varepsilon$. Then for each $y \in X$ and $z \in X$ such that $|y - x| < \delta(\varepsilon)/2$ and $|z - x| < \delta(\varepsilon)/2$, we have $|z - y| < \delta(\varepsilon)$ and $|f(z) - f(y)| < \varepsilon$.

Let C_ε^- (resp. C_ε^+) be the set of elements $\alpha \in R_0$ such that $\alpha < f(y) - \varepsilon$ (resp. $\alpha > f(y) + \varepsilon$) for some $y \in X$ satisfying $|y - x| < \delta(\varepsilon)/2$. Let C^- (resp. C^+) be the union of subsets of the form C_ε^- (resp. C_ε^+) for $\varepsilon \in R_0^{>0}$.

Claim 1: *for each $a \in C^-$ and each $b \in C^+$, we have $a < b$. In particular, the set $C^- \cap C^+$ is empty.*

There exist y_1 and y_2 in X , and ε_1 and ε_2 in $R_0^{>0}$, such that $|y_1 - x| < \delta(\varepsilon_1)/2$, $|y_2 - x| < \delta(\varepsilon_2)/2$, $a < f(y_1) - \varepsilon_1$ and $b > f(y_2) + \varepsilon_2$. Let $\delta = \min\{\delta(\varepsilon_1), \delta(\varepsilon_2)\}$. We fix $y \in X$ such that $|y - x| < \delta/2$. Then we have $|f(y) - f(y_1)| < \varepsilon_1 < f(y_1) - a$ and $|f(y) - f(y_2)| < \varepsilon_2 < b - f(y_2)$, so we obtain $a < f(y) < b$.

Claim 2: *for each $r \in R_0^{>0}$, there is $a \in C^-$ and $b \in C^+$ such that $b - a < r$.*

Let $y \in X$ such that $|y - x| < \delta(r/3)/2$. Then for $a = f(y) - r/2$ and $b = f(y) + r/2$, we have $a \in C_{r/3}^-$ and $b \in C_{r/3}^+$, and we obtain $a \in C^-$, $b \in C^+$ and $b - a < r$.

Conclusion: By [3, §2, Lemma 2.8] (or [2]), there is a unique $\omega \in R$ satisfying $a < \omega < b$ for every $a \in C^-$ and $b \in C^+$. Hence for each $\varepsilon \in R_0^{>0}$, there is $\delta(= \delta(\varepsilon)) \in R_0^{>0}$ such that whenever $|y - x| < \delta/2$ for $y \in X$, we have $\omega \geq f(y) - \varepsilon$ and $\omega \leq f(y) + \varepsilon$, so $|f(y) - \omega| \leq \varepsilon$. Now $\omega = \lim_{\substack{y \in X \\ y \rightarrow x}} f(y)$ exists. \square

Lemma 2.4.. – *Let k be an integer. If E and F are two closed definable subsets of R_0^k , then $\overline{E \cap F} = \overline{E} \cap \overline{F}$. In particular, if $\overline{E} \cap \overline{F}$ is non-empty, then $E \cap F$ is non-empty too.*

PROOF – We have just to prove that $\overline{E \cap F}$ contains $\overline{E} \cap \overline{F}$. Let $x \in \overline{E} \cap \overline{F}$. For each $\varepsilon \in R_0^{>0}$, we fix $u_\varepsilon \in E$ such that $|u_\varepsilon - x| < \varepsilon$. Let $B_\varepsilon = \{y \in R_0^k \mid |u_\varepsilon - y| \leq \varepsilon\}$. Since $x \in \overline{E} \cap \overline{F}$ and since $|u_\varepsilon - x| < \varepsilon$, we have $x \in \overline{E_\varepsilon} \cap \overline{F_\varepsilon}$ where $E_\varepsilon = B_\varepsilon \cap E$ and $F_\varepsilon = B_\varepsilon \cap F$. Moreover, we note that E_ε and F_ε are closed and bounded definable subsets of R_0^k .

We show that $E_\varepsilon \cap F_\varepsilon$ is non-empty. Let $f_\varepsilon : E_\varepsilon \times F_\varepsilon \rightarrow R_0$ defined by $f_\varepsilon(z, z') = |z - z'|$. It is a definable continuous function, so its image is closed and bounded (see [4, Chapter 6 §1]). For each $\eta \in R_0^{>0}$, there exist $u \in E_\varepsilon$ and $v \in F_\varepsilon$ such that $|u - v| < \eta$, so we have $f_\varepsilon(u, v) = |u - v| < \eta$. Since the image of f_ε is closed and bounded, this implies that it contains zero. Hence there exist $a \in E_\varepsilon$ and $b \in F_\varepsilon$ such that $f_\varepsilon(a, b) = 0$. Now we have $a = b \in E_\varepsilon \cap F_\varepsilon$, and $E_\varepsilon \cap F_\varepsilon$ is non-empty.

Since $|u_\varepsilon - x| < \varepsilon$, we have $|y - x| < 2\varepsilon$ for any $y \in B_\varepsilon$. Thus, the previous paragraph proves that for each $\varepsilon \in R_0^{>0}$, there exists $y \in E \cap F$ such that $|y - x| \leq 2\varepsilon$, so $x \in \overline{E \cap F}$, as desired. \square

Corollary 2.5.. – *For each subset X of R_0^k , and each $x \in \check{X}$, there is a closed and bounded definable subset F of X such that $x \in \overline{F}$.*

PROOF – Since $x \in \check{X}$, there exists $x_0 \in X$ such that $|x - x_0| < 1$. We consider $F_1 = \{y \in R_0^k \mid |y - x_0| \leq 1\}$. Then F_1 is a closed and bounded definable subset of R_0^k . By density of R_0 in R and since $|x - x_0| < 1$, we have $x \in \overline{F_1}$.

Moreover, by the definition of \check{X} , there is a closed definable subset F_2 of X such that $x \in \overline{F_2}$. Then $F = F_1 \cap F_2$ is a closed and bounded definable subset of X , and Lemma 2.4 gives $x \in \overline{F}$. \square

Proposition 2.6.. – *Let k be an integer. If $\{A_1, \dots, A_m\}$ is a partition of R_0^k into definable subsets, then $\{\check{A}_1, \dots, \check{A}_m\}$ is a partition of R^k .*

PROOF – First we show that R^k is the union of $\check{A}_1, \dots, \check{A}_m$. Since each definable subset of R_0^k has a decomposition into cells (see [4, Chapter 3 §2] for more details), we may assume that A_1, \dots, A_m are cells.

Let $x \in R^k$. We show that $x \in \check{A}_j$ for some $j \in \{1, \dots, m\}$. We may assume $x \notin R_0^k$. By finiteness of the partition $\{A_1, \dots, A_m\}$, there exists $r \in R^{>0}$ such that, for any $s \in]0, r[$, the set $I = \{i \in \{1, \dots, m\} \mid \exists a \in A_i, |x - a| < s\}$ is constant. Since R_0 is dense in R , the set I is non-empty, and by the definition of I , the point x is in the R^k -closure $\overline{A_i}$ of A_i for each $i \in I$.

Let d be the smallest integer such that there is $j \in \{1, \dots, m\}$ and a definable subset B of A_j of dimension d with x contained in \overline{B} . Let $B_i = A_i \cap \partial B$ for each $i \in \{1, \dots, m\}$. Then for each $i \in \{1, \dots, m\}$, the subset B_i of A_i is definable (see [4, Chapter 1 §3]) and we have $\dim B_i \leq \dim \partial B < \dim B = d$ [4, Chapter 4 §1]. By the minimality of d , the point x is contained in the R^k -closure $\overline{B_i}$ of B_i for no $i \in \{1, \dots, m\}$. Consequently, there is $t \in R^{>0}$ such that $|y - x| > t$ for any $y \in \cup_{i=1}^m B_i$, and we may choose $t \in R_0$ as R_0 is dense in R . Since $\cup_{i=1}^m B_i = \partial B$ and since $x \in \overline{B}$, there exists $b_0 \in B$ such that $|b_0 - x| < t/2$. Let $B_f = \{b \in B \mid |b - b_0| \leq t/2\}$. By the choices of t and b_0 , we have $x \in \overline{B_f}$ and $\partial B \cap B_f = \emptyset$. This implies that the set B_f is a closed definable subset of B and that x belongs to $\overset{\circ}{A}_j$, as desired.

We show that $\overset{\circ}{A}_i \cap \overset{\circ}{A}_j = \emptyset$ for any distinct elements i and j of $\{1, \dots, m\}$. Otherwise there is a closed definable subset F_i (resp. F_j) of A_i (resp. A_j) such that $\overline{F_i} \cap \overline{F_j} \neq \emptyset$. By Lemma 2.4, the set $F_i \cap F_j$ is non-empty, contradicting $A_i \cap A_j = \emptyset$. Thus $\{\overset{\circ}{A}_1, \dots, \overset{\circ}{A}_m\}$ is a partition of R^k . \square

3. PROOF OF THE MAIN THEOREM

We provide three preparatory results before the final argument.

Lemma 3.1. – *Let $S_0 = \{(u, v) \in R_0^2 \mid u < v\}$ and $S = \{(u, v) \in R^2 \mid u < v\}$. Then we have $S = \overset{\circ}{S}_0$.*

PROOF – First we show that $\overset{\circ}{S}_0$ contains S . Let $(u, v) \in S$. Then there exists $r \in R_0^{>0}$ such that $v - u > r$. Let $(u_0, v_0) \in R_0^2$ such that $|u - u_0| < r/4$ and $|v - v_0| < r/4$, and let $F = \{(x, y) \in R_0^2 \mid |x - u_0| \leq r/4, |y - v_0| \leq r/4\}$. In particular, we have $(u, v) \in \overline{F}$. Moreover we have $v_0 - u_0 > r/2$, so we obtain $y - x > 0$ for each $(x, y) \in F$, and F is a closed definable subset of S_0 . Consequently (u, v) belongs to $\overset{\circ}{S}_0$, and $\overset{\circ}{S}_0$ contains S .

Now we show that S contains $\overset{\circ}{S}_0$. Let $(u, v) \in \overset{\circ}{S}_0$. By Corollary 2.5, there is a closed and bounded definable subset F_0 of S_0 such that $(u, v) \in \overline{F_0}$. Let $f : F_0 \rightarrow R_0$ defined by $f(x, y) = y - x$. Then f is a definable continuous function, so its image is closed and bounded. Let $m = \min\{f(x, y) \mid (x, y) \in F_0\}$. Then we have $m > 0$ for each $(a, b) \in \overline{F_0}$. Since $F_0 \subseteq S_0$, we have $m > 0$ and we obtain $(u, v) \in S$. \square

Corollary 3.2. – *Let $T_0 = \{(u, v) \in R_0^2 \mid u = v\}$ and $T = \{(u, v) \in R^2 \mid u = v\}$. Then we have $T = \overset{\circ}{T}_0$.*

PROOF – Let $S_0 = \{(u, v) \in R_0^2 \mid u < v\}$ and $S_1 = \{(u, v) \in R_0^2 \mid u > v\}$. Then $\{S_0, S_1, T_0\}$ is a partition of R_0^2 , and Proposition 2.6 says that $\{\overset{\circ}{S}_0, \overset{\circ}{S}_1, \overset{\circ}{T}_0\}$ is a partition of R^2 . Now Lemma 3.1 gives $\overset{\circ}{T}_0 = R^2 \setminus (\overset{\circ}{S}_0 \cup \overset{\circ}{S}_1) = T$. \square

Lemma 3.3. – *If G_0 (resp. H_0, K_0) denotes the graph of \cdot (resp. $+$, $-$) in R_0^3 (resp. R_0^3, R_0^2), then the graph of \cdot (resp. $+$, $-$) in R^3 (resp. R^3, R^2) is $\overset{\circ}{G}_0$ (resp. $\overset{\circ}{H}_0, \overset{\circ}{K}_0$).*

PROOF – Since \cdot is a continuous map over R_0 , its graph G_0 in R_0^3 is closed, and we have $\overset{\circ}{G}_0 = \overline{G_0}$. Moreover, since \cdot is a continuous map over R , its graph G in R^3 is closed, and since G contains G_0 , it contains $\overline{G_0}$ too. But R_0 is dense in R , hence for each $(x, y) \in R^2$ we have $(x, y, x \cdot y) \in \overline{G_0}$, and G is contained in $\overline{G_0}$. We conclude that $G = \overline{G_0} = \overset{\circ}{G}_0$, as desired.

In the same way, we show that the graph of $+$ (resp. $-$) in R^3 (resp. R^2) is \check{H}_0 (resp. \check{K}_0). \square

PROOF OF THEOREM 1.2 – We denote by \mathcal{L}_1 the language of $\mathcal{M}_1 = (R_0, <, \dots)$. For each function symbol f of \mathcal{L}_1 with arity k , we consider a relation symbol S_f such that $S_f^{\mathcal{M}_1}$ is the graph of $f^{\mathcal{M}_1}$, and for each constant symbol c of \mathcal{L}_1 , we consider a relation symbol S_c such that $S_c^{\mathcal{M}_1} = c^{\mathcal{M}_1}$. We obtain a relational language \mathcal{L} and a structure $\mathcal{M}_0 = (R_0, <, \dots)$ in \mathcal{L} . We have just to prove that there is a unique \mathcal{L} -structure \mathcal{M} with domain R such that \mathcal{M} is an elementary extension of \mathcal{M}_0 .

We note that, for any integer k , a subset X of R_0^k is definable (in \mathcal{M}_1) if and only if it is definable in \mathcal{M}_0 .

Uniqueness:

First we assume that the structure \mathcal{M} exists. Let S be any relation symbol of arity k of \mathcal{L} . Let F be a closed and bounded definable subset of $S^{\mathcal{M}_0}$, and let $\varphi(\bar{x}, \bar{a})$ be an \mathcal{L} -formula with free variables $\bar{x} = (x_1, \dots, x_k)$ and parameters $\bar{a} \in R_0^m$ such that F is defined by $\varphi(\bar{x}, \bar{a})$. Let \tilde{F} be the subset of R^k defined by $\varphi(\bar{x}, \bar{a})$. Since \mathcal{M} is an elementary extension of \mathcal{M}_0 , then \tilde{F} contains F and it is closed and bounded in R^k . Thus \tilde{F} contains \overline{F} . By Corollary 2.5, this implies that $S^{\mathcal{M}}$ contains $\check{S}^{\mathcal{M}_0}$.

In the same way, the complementary of $S^{\mathcal{M}}$ in R^k contains $\widehat{R_0^k \setminus S^{\mathcal{M}_0}}$. Now Proposition 2.6 gives $S^{\mathcal{M}} = \check{S}^{\mathcal{M}_0}$, so, if it exists such a structure \mathcal{M} , then it is unique.

Existence:

We consider the \mathcal{L} -structure $\mathcal{M} = (R, <, \dots)$ where for each relation symbol S of arity k of \mathcal{L} , we define $S^{\mathcal{M}}$ by $S^{\mathcal{M}} = \check{S}^{\mathcal{M}_0}$. By Lemmas 3.1 and 3.3 and Corollary 3.2, it is sufficient to show that \mathcal{M} is an elementary extension of \mathcal{M}_0 .

First we note that for each relation symbol S of arity k of \mathcal{L} , we have $S^{\mathcal{M}_0} = \check{S}^{\mathcal{M}_0} \cap R_0^k$ (Remark 2.2), so \mathcal{M}_0 is a substructure of \mathcal{M} .

Claim 1: *if A is a definable subset of R_0^k for an integer k , then we have $\check{A}_0 = R^k \setminus \check{A}$ where $A_0 = R_0^k \setminus A$.*

Since $\{A, A_0\}$ is a partition of R_0^k , then Proposition 2.6 says that $\{\check{A}, \check{A}_0\}$ is a partition of R^k .

Claim 2: *if A and B are two definable subsets of R_0^k for an integer k , then we have $\check{A} \cap \check{B} = \widehat{A \cap B}$.*

It is sufficient to prove that $\widehat{A \cap B}$ contains $\check{A} \cap \check{B}$. Let $x \in \check{A} \cap \check{B}$. Then there exist a closed definable subset E of A and a closed definable subset F of B such that $x \in \overline{E} \cap \overline{F}$. Now x belongs to $\overline{E \cap F} \subseteq \widehat{A \cap B}$ by Lemma 2.4, and we obtain $\check{A} \cap \check{B} = \widehat{A \cap B}$.

Claim 3: *for any two integers k and l , if A and B are definable subsets of R_0^k and R_0^l respectively, then we have $\check{A} \times \check{B} = \widehat{A \times B}$.*

Let $x \in \check{A} \times \check{B}$. We show that x belongs to $\widehat{A \times B}$. We have $x = (u, v)$ for $u \in \check{A}$ and $v \in \check{B}$. Then there are two closed definable subsets F and G of A and B respectively such that $u \in \overline{F}$ and $v \in \overline{G}$. For each $\varepsilon \in R^{>0}$, there $f \in F$ and

$g \in G$ such that $|u - f| < \varepsilon$ and $|v - g| < \varepsilon$, so $|(u, v) - (f, g)| < \varepsilon$. Consequently, $x = (u, v)$ belongs to $\overline{F \times G} \subseteq \overline{A \times B}$.

By Corollary 2.5, for any $x \in \overline{A \times B}$, there is a closed and bounded definable subset H of $A \times B$ such that $x \in \overline{H}$. If H_1 (resp. H_2) denotes the image of H by the projection $\pi_1 : A \times B \rightarrow A$ (resp. $\pi_2 : A \times B \rightarrow B$), then by the continuity of the projections maps, the set H_1 (resp. H_2) is closed and bounded, and H is contained in $H_1 \times H_2$. Now \overline{H} is contained in $\overline{H_1 \times H_2}$. But $\overline{H_1 \times H_2}$ is closed in R^{k+l} and it contains $H_1 \times H_2$, so it contains $\overline{H_1 \times H_2}$. Hence \overline{H} is contained in $\overline{H_1 \times H_2}$, and $\overline{A \times B}$ contains $\overline{A \times B}$.

Claim 4: let $\pi : R^{k+1} \rightarrow R^k$ be the projection on the first k coordinates for an integer k . If A is a definable subset of R_0^{k+1} , then we have $\pi(\check{A}) = \overline{\pi(\check{A})}$.

First we note that the restriction $\pi|_{R_0^{k+1}} : R_0^{k+1} \rightarrow R_0^k$ of π to R_0^{k+1} is definable and continuous. In particular, the set $\pi(A)$ is definable.

Let $x \in \check{A}$. By Corollary 2.5, there is a closed and bounded definable subset F of A such that $x \in \overline{F}$. Then, for each $\varepsilon \in R^{>0}$, there exists $y \in F$ such that $|y - x| < \varepsilon$, so we have $|\pi(y) - \pi(x)| < \varepsilon$, and thus we obtain $\pi(x) \in \overline{\pi(F)}$. But the restriction $\pi|_{R_0^{k+1}}$ is definable and continuous, so $\pi(F)$ is a closed and bounded definable subset contained in $\pi(A)$. Hence $\pi(x)$ belongs to $\overline{\pi(A)}$, and $\overline{\pi(\check{A})}$ contains $\pi(\check{A})$.

Let $x \in \overline{\pi(\check{A})}$. We show that x belongs to $\pi(\check{A})$. By definable choice [4, Chapter 6 §1], there is a definable map $f : \pi(A) \rightarrow R_0$ such that $\{(\alpha, f(\alpha)) \mid \alpha \in \pi(A)\}$ is contained in A . By cell decomposition [4, Chapter 3 §2], there are finitely many cells C_1, \dots, C_s of $\pi(A)$ such that $\pi(A) = \cup_{i=1}^s C_i$ and f is continuous on C_i for each i . By Proposition 2.6, we have $x \in \check{C}_r$ for some $r \in \{1, \dots, s\}$. By Corollary 2.5, there is a closed and bounded definable subset G of C_r such that $x \in \overline{G}$. But f is continuous on G , so the graph Γ of its restriction $f|_G : G \rightarrow R_0$ to G is a closed and definable subset of A . Moreover, the continuity of f on G implies its uniform continuity on G [4, Chapter 6 §1], hence the following limit exists by Lemma 2.3:

$$u = \lim_{\substack{y \in G \\ y \rightarrow x}} f(y)$$

Now (x, u) belongs to $\overline{\Gamma}$, and since Γ is a closed and definable subset of A , we obtain $(x, u) \in \check{A}$ and $x \in \pi(\check{A})$.

Claim 5: if $\varphi(\overline{x}, \overline{a})$ is an atomic formula with free variables $\overline{x} = (x_1, \dots, x_k)$ and parameters $\overline{a} = (a_1, \dots, a_m)$ in R_0^m , and if A is the definable subset of R_0^k defined by $\varphi(\overline{x}, \overline{a})$, then \check{A} is the \mathcal{M} -definable subset of R^k defined by $\varphi(\overline{x}, \overline{a})$.

Let S be a relation symbol such that $\varphi(\overline{x}, \overline{a}) = S(\overline{x}, \overline{a})$. By the definition of $S^{\mathcal{M}}$, we have $S^{\mathcal{M}} = \check{S}^{\mathcal{M}_0}$. Therefore, if $B = S^{\mathcal{M}_0}$ is the subset of R_0^{k+m} defined by $\varphi(\overline{x}, \overline{y})$, if $C = R_0^k \times \{\overline{a}\}$, and if $\pi : R^{k+m} \rightarrow R^k$ is the projection on the first k coordinates, then we have $A = \pi(B \cap C)$. In the same way, the \mathcal{M} -definable subset of R^k defined by $\varphi(\overline{x}, \overline{a})$ is $\pi(S^{\mathcal{M}} \cap (R^k \times \{\overline{a}\}))$. Since by Claim 3 we have $\check{C} = \check{R}_0^k \times \overline{\{\overline{a}\}} = R^k \times \{\overline{a}\}$, and since by Claims 2 and 4 we have $\check{A} = \pi(\check{B} \cap \check{C})$, Claim 5 is proven.

Claim 6: let $\varphi(\bar{x}, y, \bar{a})$ be a formula with free variables $\bar{x} = (x_1, \dots, x_k)$ and y , and parameters $\bar{a} \in R_0^m$. Let A be the subset of R_0^{k+1} defined by $\varphi(\bar{x}, y, \bar{a})$, and B be the subset of R_0^k defined by $\exists y \varphi(\bar{x}, y, \bar{a})$. If the subset of R^{k+1} defined by $\varphi(\bar{x}, y, \bar{a})$ is \check{A} , then the subset of R^k defined by $\exists y \varphi(\bar{x}, y, \bar{a})$ is \check{B} .

This follows from Claim 4.

Claim 7: let $\varphi(\bar{x}, \bar{a})$ be a formula with free variables $\bar{x} = (x_1, \dots, x_k)$ and parameters $\bar{a} \in R_0^m$. Let A be the subset of R_0^k defined by $\varphi(\bar{x}, \bar{a})$, and B be the subset of R_0^k defined by $\neg\varphi(\bar{x}, \bar{a})$. If the subset of R^k defined by $\varphi(\bar{x}, \bar{a})$ is \check{A} , then the subset of R^k defined by $\neg\varphi(\bar{x}, \bar{a})$ is \check{B} .

This follows from Claim 1.

Claim 8: let $\varphi(\bar{x}, \bar{a})$ and $\phi(\bar{x}, \bar{a})$ be formulas with free variables $\bar{x} = (x_1, \dots, x_k)$ and parameters $\bar{a} \in R_0^m$. Let A be the subset of R_0^k defined by $\varphi(\bar{x}, \bar{a})$, and B be the subset of R_0^k defined by $\phi(\bar{x}, \bar{a})$. If the subset of R^k defined by $\varphi(\bar{x}, \bar{a})$ is \check{A} and the one defined by $\phi(\bar{x}, \bar{a})$ is \check{B} , then the subset of R^k defined by $\varphi(\bar{x}, \bar{a}) \wedge \phi(\bar{x}, \bar{a})$ is $\check{A \cap B}$.

This follows from Claim 2.

Conclusion: it follows from Claims 5, 6, 7 and 8, and from Remark 2.2, that the structure \mathcal{M} is an elementary extension of \mathcal{M}_0 . \square

REFERENCES

- [1] Antongiulio Fornasiero. Tame structures and open cores. 2010. <http://www.logique.jussieu.fr/modnet/Publications/Preprint%20server/papers/237/237.pdf>.
- [2] Dana Scott. On completing ordered fields. In *Applications of Model Theory to Algebra, Analysis, and Probability (Internat. Sympos., Pasadena, Calif., 1967)*, pages 274–278. Holt, Rinehart and Winston, New York, 1969.
- [3] Saharon Shelah. Quite complete real closed fields. *Israel J. Math.*, 142:261–272, 2004.
- [4] Lou van den Dries. *Tame topology and o-minimal structures*, volume 248 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 1998.

LABORATOIRE DE MATHÉMATIQUES ET APPLICATIONS, UNIVERSITÉ DE POITIERS
E-mail address: olivier.frecon@math.univ-poitiers.fr